



R. Scalzo, G. Aldering, B. C. Lee, S. Loken, P. Nugent, S. Perlmutter, J. Siegrist, R.C. Thomas, L. Wang, W. M. Wood-Vasey (Lawrence Berkeley National Lab),
G. Adam, R. Bacon, C. Bonnaud, L. Capoani, D. Dubet, F. Henault, B. Lantz, J.-P. Lemonnier, A. Pecontal, E. Pecontal (Centre de Recherche Astronomique de Lyon),
N. Blanc, G. Boudoul, S. Bongard, A. Castera, Y. Copin, E. Gangler, G. Smadja (Institut de Physique Nucleaire de Lyon),
R. Kessler (Kavli Institute for Cosmological Physics, University of Chicago), P. Antilogus, P. Astier, E. Berrelet, G. Garavini,
S. Gilles, L.-A. Guevara, D. Imbault, C. Juramy, R. Pain, R. Taillet, D. Vincent (Laboratoire de Physique Nucleaire et de Haute Energies de Paris),
C. Baltay, A. Bauer, D. Rabinovitz (Yale University)



The Search:



A large yellow industrial machine, likely a wind turbine component, is shown. A red bounding box highlights a specific mechanical joint or assembly on the left side of the machine.



SN2004gc

Red channel lens array

1 spectra per lens
=225 spectra per channel

15x15 lenses
(1 lens = $0.4'' \times 0.4''$)

Sum the lenses/spectra with light from the SN

x 2 channels
(blue + red channels)

SNfactory PRELIMINARY

Unfiltered flux (Arbitrary unit)

Wavelength (microm)

In order to obtain spectrophotometric observations of ~ 300 SNe Ia, the SNfactory constructed the SuperNova Integral Field Spectrograph (SNIFS), which was installed on the UH 2.2-m telescope on Mauna Kea in April 2004. At present 20% of all observing time on the UH 2.2-m is devoted to observing supernovae with SNIFS. The instrument features a dual-channel optical spectrograph (left) and an imaging/guiding camera (right).

Each spectrographic channel exploits a 15x15 microlens array to image 225 spectra across the 6"x6" field of view.

During each spectroscopic exposure, the imaging camera makes multi-band photometric observations of stars in an adjacent field using custom-designed filters. By comparing the observed brightnesses of these stars to the brightnesses observed on a photometric night, the corresponding supernova spectrum can be corrected for atmospheric extinction even on non-photometric nights.

725 nm

950 nm

850 nm

590 nm

V
550 nm

400 nm

For more information on the SNfactory, contact Richard Scalzo (RAScalzo@lbl.gov) or Greg Aldering (GAldering@LBL.gov) or visit out webpage at <http://snfactory.lbl.gov/>